

Color

- Reading:
 - Chapter 6, Forsyth & Ponce
- Optional reading:
 - Chapter 4 of Wandell, Foundations of Vision, Sinauer, 1995 has a good treatment of this.

Feb. 19, 2004
MIT 6.891
Prof. Freeman for Prof. Darrell

Why does a visual system need color?



<http://www.hobbyfinc.com/grpl/p15019.jpg>


Why does a visual system need color? (an incomplete list...)

- To tell what food is edible.
- To distinguish material changes from shading changes.
- To group parts of one object together in a scene.
- To find people's skin.
- Check whether a person's appearance looks normal/healthy.
- To compress images

Lecture outline


- Color physics.
- Color perception and color matching..

Color




4.1 NEWTON'S SUMMARY DRAWING of his experiments with light. Using a point source of light and a prism, Newton separated sunlight into its fundamental components. By reconverging the rays, he also showed that the decomposition is reversible.
From Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995

Spectral colors



<http://hyperphysics.phy-astr.gsu.edu/hbase/vision/speccol.html#c2>

Radiometry (review)



Horn, 1986
Figure 10-7. The bidirectional reflectance distribution function is the ratio of the radiance of the surface patch as viewed from the direction (θ_e, ϕ_e) to the irradiance resulting from illumination from the direction (θ_i, ϕ_i) .

radiance


$$BRDF = f(\theta_i, \phi_i, \theta_e, \phi_e) = \frac{L(\theta_e, \phi_e)}{E(\theta_i, \phi_i)}$$

irradiance

Radiometry for colour

- All definitions are now “per unit wavelength”
- All units are now “per unit wavelength”
- All terms are now “spectral”
- Radiance becomes spectral radiance
 - watts per square meter per steradian per unit wavelength
- Irradiance becomes spectral irradiance
 - watts per square meter per unit wavelength

Radiometry for color



Horn, 1986
Figure 10-7. The bidirectional reflectance distribution function is the ratio of the radiance of the surface patch as viewed from the direction (θ_e, ϕ_e) to the irradiance resulting from illumination from the direction (θ_i, ϕ_i) .

Spectral radiance


$$BRDF = f(\theta_i, \phi_i, \theta_e, \phi_e, \lambda) = \frac{L(\theta_e, \phi_e, \lambda)}{E(\theta_i, \phi_i, \lambda)}$$

Spectral irradiance

Simplified rendering models: reflectance




Often are more interested in relative spectral composition than in overall intensity, so the spectral BRDF computation simplifies a wavelength-by-wavelength multiplication of relative energies.



Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995


Simplified rendering models: transmittance



Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995

How measure those spectra: Spectrophotometer


(just like Newton's diagram...)



4.2 A SPECTRORADIOMETER is used to measure the spectral power distribution of light. (A) A schematic design of a spectroradiometer includes a means for separating the input light into its different wavelengths and a detector for measuring the energy at each of the separate wavelengths. (B) The color names associated with the appearance of lights at a variety of wavelengths are shown. After Wyszecki and Stiles, 1982.


Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995

Two illumination spectra





Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995


Some reflectance spectra



Color names for cartoon spectra



Additive color mixing




When colors combine by *adding* the color spectra. Examples that follow this mixing rule: CRT phosphors, multiple projectors aimed at a screen, Polachrome slide film.

Red and green make...

Yellow!

Subtractive color mixing



When colors combine by *multiplying* the color spectra. Examples that follow this mixing rule: most photographic films, paint, cascaded optical filters, crayons.

Cyan and yellow (in crayons, called "blue" and yellow) make...

Green!

demos

- Additive color
- Subtractive color


Low-dimensional models for color spectra

$$e(\lambda) = \begin{pmatrix} \vdots \\ \vdots \\ \vdots \end{pmatrix} = \begin{pmatrix} \vdots & \vdots & \vdots \\ E_1(\lambda) & E_2(\lambda) & E_3(\lambda) \\ \vdots & \vdots & \vdots \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \\ \omega_3 \end{pmatrix}$$

How to find a linear model for color spectra:

- form a matrix, D, of measured spectra, 1 spectrum per column.
- [u, s, v] = svd(D) satisfies $D = u * s * v'$
- the first n columns of u give the best (least-squares optimal) n-dimensional linear bases for the data, D:
 $D \approx u(:, 1:n) * s(1:n, 1:n) * v(1:n, :)$


Basis functions for Macbeth color checker



9.9 BASIS FUNCTIONS OF THE LINEAR MODEL FOR THE MACBETH COLORCHECKER. The surface-reflection functions in the collection vary smoothly with wavelength, as do the basis functions. The first basis function is all positive and explains the most variance in the surface-reflection functions. The basis functions are ordered in terms of their relative significance for reducing the error in the linear-model approximation to the surfaces.

Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995

n-dimensional linear models for color spectra



9.8 A LINEAR MODEL TO APPROXIMATE THE SURFACE REFLECTANCES IN THE MACBETH COLORCHECKER. The panels in each row of this figure show the surface-reflection functions of six colored surfaces (shaded lines) and the approximation to these functions using a linear model (solid lines). The approximations using linear models with (A) three, (B) two, and (C) one dimension are shown.

Forsyth & Ponce


Outline

- Color physics.
- Color perception and color matching.

Why specify color numerically?





- Accurate color reproduction is commercially valuable
 - Many products are identified by color ("golden" arches);
- Few color names are widely recognized by English speakers
 - About 10; other languages have fewer/more, but not many more.
 - It's common to disagree on appropriate color names.
- Color reproduction problems increased by prevalence of digital imaging - eg. digital libraries of art.
 - How do we ensure that everyone sees the same color?

Color standards are important in industry




An assumption that sneaks in here


- We know color appearance really depends on:
 - The illumination
 - Your eye's adaptation level
 - The colors and scene interpretation surrounding the observed color.
- But for now we will assume that the spectrum of the light arriving at your eye completely determines the perceived color.




Color matching experiment 1




Color matching experiment 2



Color matching experiment 2



Color matching experiment 2






Color matching experiment 2

We say a "negative" amount of p_2 was needed to make the match, because we added it to the test color's side.



The primary color amounts needed for a match:



4.12 THE COLOR-MATCHING EXPERIMENT SATISFIES THE PRINCIPLE OF SUPERPOSITION. In parts (A) and (B), test lights are matched by a mixture of three primary lights. In part (C) the sum of the test lights is matched by the additive mixture of the primaries, demonstrating superposition.

Grassman's Laws

- For color matches:
 - symmetry: $U=V \Leftrightarrow V=U$
 - transitivity: $U=V$ and $V=W \Rightarrow U=W$
 - proportionality: $U=V \Leftrightarrow tU=tV$
 - additivity: if any two (or more) of the statements $U=V$, $W=X$, $(U+W)=(V+X)$ are true, then so is the third
- These statements are as true as any biological law. They mean that additive color matching is linear.

Forsyth & Ponce


Measure color by color-matching paradigm

- Pick a set of 3 primary color lights.
- Find the amounts of each primary, e_1, e_2, e_3 , needed to match some spectral signal, t .
- Those amounts, e_1, e_2, e_3 , describe the color of t . If you have some other spectral signal, s , and s matches t perceptually, then e_1, e_2, e_3 will also match s .
- Why this is useful—it lets us:
 - Predict the color of a new spectral signal
 - Translate to representations using other primary lights.

How to do this, mathematically

- Pick a set of primaries, $p_1(\lambda), p_2(\lambda), p_3(\lambda)$
- Measure the amount of each primary, $c_1(\lambda), c_2(\lambda), c_3(\lambda)$ needed to match a monochromatic light, $t(\lambda)$ at each spectral wavelength λ (pick some spectral step size).


Color matching functions for a particular set of monochromatic primaries



Using the color matching functions to predict the primary match to a new spectral signal

Store the color matching functions in the rows of the matrix, C

$$C = \begin{pmatrix} c_1(\lambda_1) & \dots & c_1(\lambda_N) \\ c_2(\lambda_1) & \dots & c_2(\lambda_N) \\ c_3(\lambda_1) & \dots & c_3(\lambda_N) \end{pmatrix}$$



Let the new spectral signal to be characterized by the vector t .

$$\vec{t} = \begin{pmatrix} t(\lambda_1) \\ \vdots \\ t(\lambda_N) \end{pmatrix}$$

Then the amounts of each primary needed to match t are:

$$C\vec{t}$$

How do you translate colors between different systems of primaries?


$$\begin{aligned} \text{Red } p_1 &= (0.0 \ 0.0 \ 0.0 \ \dots \ 0.1 \ 0)^T & \text{Orange } p'_1 &= (0.0 \ 0.2 \ 0.3 \ 4.5 \ 7 \ \dots \ 2.1)^T \\ \text{Green } p_2 &= (0.0 \ \dots \ 0.1 \ 0 \ \dots \ 0)^T & \text{Yellow } p'_2 &= (0.1 \ 0.44 \ 2.1 \ \dots \ 0.3 \ 0)^T \\ \text{Blue } p_3 &= (0.1 \ 0.0 \ \dots \ 0.0 \ 0)^T & \text{Purple } p'_3 &= (1.2 \ 1.7 \ 1.6 \ \dots \ 0.0)^T \end{aligned}$$

Primary spectra, P

Color matching functions, C


Primary spectra, P'

Color matching functions, C'



So color matching functions translate like this:


From previous slide $C\vec{t} = CP' C'\vec{t}$ But this holds for any input spectrum, t , so...



$$C = \underbrace{CP' C'}_{\text{a 3x3 matrix}}$$

P' are the old primaries
C are the new primaries' color matching functions

How do you translate from the color in one set of primaries to that in another?



$$e = \underbrace{CP'}_{\text{the same 3x3 matrix}} e'$$

P' are the old primaries
C are the new primaries' color matching functions

What's the machinery in the eye?


Eye Photoreceptor responses




(Where do you think the light comes in?)

The intricate layers and connections of nerve cells in the retina were drawn by the famed Spanish anatomist Santiago Ramón y Cajal around 1900. Rod and cone cells are at the top. Optic nerve fibers leading to the brain may be seen at bottom right.

Human Photoreceptors




Human eye photoreceptor spectral sensitivities




3.3 SPECTRAL SENSITIVITIES OF THE L-, M-, AND S-CONES in the human eye. The measurements are based on a light source at the cornea, so that the wavelength loss due to the cornea, lens, and other inert pigments of the eye plays a role in determining the sensitivity. Source: Stockman and MacLeod, 1993.

Are the color matching functions we observe obtainable from some 3x3 matrix transformation of the human photopigment response curves?

Color matching functions (for a particular set of spectral primaries)



Comparison of color matching functions with best 3x3 transformation of cone responses



4.20 COMPARISON OF CONE PHOTOCURRENT RESPONSES AND THE COLOR-MATCHING FUNCTIONS. The cone photocurrent spectral responsivities are within a linear transformation of the color-matching functions, after a correction has been made for the optics and inert pigments in the eye. The smooth curves show the Stiles and Burch (1959) color-matching functions. The symbols show the matches predicted from the photocurrents of the three types of macaque cones. The predictions included a correction for absorption by the lens and other inert pigments in the eye. Source: Baylor, 1987.


Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995

Since we can define colors using almost any set of primary colors, let's agree on a set of primaries and color matching functions for the world to use...

CIE XYZ color space

- Commission Internationale d'Eclairage, 1931
- "...as with any standards decision, there are some irritating aspects of the XYZ color-matching functions as well...no set of physically realizable primary lights that by direct measurement will yield the color matching functions."
- "Although they have served quite well as a technical standard, and are understood by the mandarins of vision science, they have served quite poorly as tools for explaining the discipline to new students and colleagues outside the field."


Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995





4.14 THE XYZ STANDARD COLOR-MATCHING FUNCTIONS. In 1931 the CIE standardized a set of color-matching functions for image interchange. These color-matching functions are called $\bar{x}(\lambda)$, $\bar{y}(\lambda)$, and $\bar{z}(\lambda)$. Industrial applications commonly describe the color properties of a light source using the three primary intensities needed to match the light source that can be computed from the XYZ color-matching functions.

CIE XYZ: Color matching functions are positive everywhere, but primaries are imaginary. Usually draw x, y, z , where $x = X/(X+Y+Z)$
 $y = Y/(X+Y+Z)$

Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995



A qualitative rendering of the CIE (x,y) space. The blobby region represents visible colors. There are sets of (x, y) coordinates that don't represent real colors, because the primaries are not real lights (so that the color matching functions could be positive everywhere).




A plot of the CIE (x,y) space. We show the spectral locus (the colors of monochromatic lights) and the black-body locus (the colors of heated black-bodies). I have also plotted the range of typical incandescent lighting.

Some other color spaces...


Uniform color spaces

- McAdam ellipses (next slide) demonstrate that differences in x,y are a poor guide to differences in color
- Construct color spaces so that differences in coordinates are a good guide to differences in color.

Forsyth & Ponce



HSV hexcone



Forsyth & Ponce

Color metamerism

Two spectra, t and s , perceptually match when


$$\vec{C}t = \vec{C}s$$

where C are the color matching functions for some set of primaries.

Graphically,

$$\begin{matrix} \boxed{C} \\ | \\ \vec{t} \end{matrix} = \begin{matrix} \boxed{C} \\ | \\ \vec{s} \end{matrix}$$

Metameric lights



4.11 METAMERIC LIGHTS. Two lights with these spectral power distributions appear identical to most observers and are called metamers. (A) An approximation to the spectral power distribution of a tungsten bulb. (B) The spectral power distribution of light emitted from a conventional television monitor whose three phosphor intensities were set to match the light in panel A in appearance.

Foundations of Vision, by Brian Wandell, Sinauer Assoc., 1995

Color constancy demo

- We assumed that the spectrum impinging on your eye determines the object color. Here's a counter-example...